

Mathematisches Forschungsinstitut Oberwolfach



Oberwolfach Preprints

OWP 2020 - 17 Neslihan Gügümcü and Sofia Lambropoulou

Braidoids

Mathematisches Forschungsinstitut Oberwolfach gGmbH Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

The MFO publishes a preprint series **Oberwolfach Preprints (OWP)**, ISSN 1864-7596, which mainly contains research results related to a longer stay in Oberwolfach, as a documentation of the research work done at the MFO. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 20 free hard copies (DIN A4, black and white copy) by surface mail.

The full copyright is left to the authors. With the submission of a manuscript, the authors warrant that they are the creators of the work, including all graphics. The authors grant the MFO a perpetual, non-exclusive right to publish it on the MFO's institutional repository.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX). Additionally, each preprint will get a Digital Object Identifier (DOI).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO) Schwarzwaldstrasse 9-11 77709 Oberwolfach-Walke Germany

 Tel
 +49 7834 979 50

 Fax
 +49 7834 979 55

 Email
 admin@mfo.de

 URL
 www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO. Copyright of the content is held by the authors.

DOI 10.14760/OWP-2020-17

1

NESLİHAN GÜGÜMCÜ AND SOFIA LAMBROPOULOU

ABSTRACT. Braidoids generalize the classical braids and form a counterpart theory to the theory of planar knotoids, just as the theory of braids does for the theory of knots. In this paper, we introduce the notion of braidoids in \mathbb{R}^2 , a closure operation for braidoids, we prove an analogue of the Alexander theorem, namely an algorithm that turns a knotoid into a braidoid, and we formulate and prove a geometric analogue of the Markov theorem for braidoids using the *L*-moves.

1. INTRODUCTION

A knotoid in an oriented surface Σ is an equivalence class of oriented open-ended knot diagrams in Σ , with two endpoints that can lie in any local region determined by the diagram. The equivalence is generated by the Reidemeister moves and isotopies of Σ , which include the swinging of an endpoint within a region free of endpoints. View Figures 1 and 2. When Σ is, in particular, the 2-sphere S^2 the knotoids are named spherical knotoids. When Σ is the plane they are named planar knotoids.

The theory of knotoids was introduced by V. Turaev in 2011 [37]. Turaev showed that the set of classical knots injects into the set of spherical knotoids, where a knot can be viewed as a knotoid with zero complexity. The complexity of a knotoid K(or height, as the term used in [17]) is the minimum number of crossings over all diagrams of K, that are created when realizing the end-to-end (underpass) closure of each diagram of K to a knot diagram. The above observation provides a strong motivation for using knotoids in computing classical knot invariants, especially those based on the number of crossings, since the computations would reduce exponentially with reducing the number of crossings in a diagram. We recall that the complete classification of knots is still a big open problem in Mathematics and it is tackled by constructing isotopy invariants for distinguishing pairs of different knots. Indeed, Turaev -among other results- defined the knot group in terms of knotoids, invariants for knotoids, such as the bracket polynomial, and he remarked on the fruitful connections of the theory with virtual knot theory. Turaev also showed that there is a surjective map from the set of planar knotoids to the set of spherical knotoids, which is not injective.

After the works [37, 16] and [7], the interest in knotoids was rekindled by the second author, who proposed the subject to the first author for her PhD study [19]. Then, in [17] the first author with L. H. Kauffman constructed new invariants for knotoids, considering the virtual end-to-end closure and employing techniques from virtual knot theory [25]. We recall that a virtual knot diagram contains classical as

¹This article is accepted to be published in Israel Journal of Mathematics

¹⁹⁹¹ Mathematics Subject Classification. 2010 Mathematics Subject Classification: 57M27; 57M25.

Key words and phrases. Braidoids, knotoids, Markov theorem, Alexander theorem.

well as virtual crossings, where a virtual crossing has no information under or over, it roughly indicates a permutation of the two acs involved. Another interesting result in [17] is the realization of a knotoid via its lifting to an open-ended curve embedded in 3-space. Through this lifting knotoids could serve as mathematical models for proteins and the theory of knotoids could be used for analyzing their topology. Consequently, in [13] invariants of spherical knotoids were applied, which rendered as much information as the use of the virtual closure, while in [14] the application of planar knotoids revealed richer structure.

In parallel, the theory of braidoids was initiated by the authors of the present paper [19, 18] for counterparting the theory of planar knotoids, just as classical braids comprise an algebraic counterpart to classical knots. A braid is a set of descending strands with paired top-to-bottom ends, and equivalence classes of braids under obvious planar isotopy and level preserving Reidemeister moves realize groups, the Artin braid groups of type A [3, 4, 24]. The paired ends of a braid can be joined to form a closure, which is a knot or a link. The inverse operation consists in turning an oriented knot or link into an isotopic closed braid and this is always possible by the classical Alexander theorem [11, 2, 9]. The Alexander theorem and its proof play a key role in the proof of the Markov theorem, which provides an equivalence relation among elements of the braid groups that corresponds precisely to the isotopy relation among oriented knots and links [34, 39, 9, 8]. The two theorems allow, in principle, for the use of braids and algebraic techniques in the study of knots and links and they were successfully used for the first time by V. F. R. Jones in his construction of the famous Jones polynomial [21, 22]. Consequently, these theorems received anew the attention of several mathematicians [35, 38, 40, 36, 31, 32, 10], whose works revealed diverse and interesting approaches.

In analogy to a braid diagram, a braidoid diagram consists in a set of descending strands such that two of them are special: one of them terminates at an endpoint, the head, and the other starts from the second endpoint, the leg. Either endpoint may lie in any region and at any height of the diagram. See Figure 5 for examples. A braidoid, then, is the equivalence class of braidoid diagrams under obvious isotopy moves analogous to the knotoid equivalence moves, as illustrated in Figures 6, 8, 9. In particular, endpoints may swing and there are also the forbidden moves in the theory, Figure 7. Hence, the notion of braidoid extends the notion of classical braid. We study braidoids in relation to knotoids by defining an appropriate closure operation, which respects the forbidden moves. The closure is defined on braidoid diagrams with labelled ends, specifying whether the joining arcs will run all over or all under the rest of the diagram.

An analogue of the classical Alexander theorem, stating that any knotoid diagram can be turned into a labeled braidoid diagram with isotopic closure, was proved in [18]. In this paper, we give a second proof of the Alexander theorem for braidoids, which is more adapted for proving an analogue of the classical Markov theorem. Indeed, we formulate and prove a geometric analogue of the Markov theorem for braidoids, using the concept of the *L*-move as introduced in [31, 32]. An *L*-move is a geometric as well as algebraic move for braids and it consists in cutting a braid arc at any point and pulling the two ends along the vertical line of the cutpoint, both over or both under the rest of the diagram, so as to obtain in the end a new pair of corresonding braid strands. The *L*-moves provide an one-move version of the classical Markov theorem [32]. The set of braidoids does not support

an obvious algebraic structure. So, it would not be possible to formulate a braidoid equivalence without the use of the L-moves. In [19, 18] a set of building blocks for braidoids is listed along with some relations that they satisfy. We then propose in [18] to encode a protein by the monomials of the building blocks for the braidoids corresponding to the knotoids related to the protein. We finally relate braidoids to classical or virtual braids by defining appropriately the underpass resp. virtual closure. We recall that a virtual braid contains classical as well as virtual crossings, see [25, 26].

For further works on knotoids and applications the interested reader may consult [1, 5, 15, 20, 28, 29].

Let us now present the organization of the paper. In Section 2 we review basics on knotoids. In Section 3 we define the notion of braidoid diagram and the notion of braidoid by introducing isotopy moves on braidoid diagrams. In Section 4, we explain a way to close a braidoid diagram with labels which will relate them to knotoids. Later, in Section 5 we describe an algorithm for turning a knotoid diagram into a labeled braidoid diagram with isotopic closure. This yields an analogue of the classical Alexander theorem for braidoids. In Section 6 we adapt the classical Lmoves, which were originally defined for braid diagrams by the second listed author in [30], for braidoid diagrams. We also introduce the fake swing moves, which along with the L-moves comprise the L-equivalence. We then prove our geometric analogue of the Markov theorem for braidoids. Finally, in Section 7 we present the underpass and virtual closures that relate a braidoid to a classical resp. a virtual braid.

2. A review of knotoids

Let [0, 1] be the unit interval and Σ be any oriented surface. A knotoid diagram K in Σ is an immersion $K : [0, 1] \to \Sigma$ that is generic in the sense that there is only a finite number of double points appearing as transversal crossings each with the extra information of under or over. The images of 0 and 1 are two distinct points disjoint from any crossings of K, too, and are called *leg* and *head* of K, respectively. Furthermore, K inherits a natural orientation from its leg to its head. In Figure 1 we show some examples of knotoid diagrams including the trivial knotoid diagram that admits no crossing.



FIGURE 1. Knotoid diagrams

Definition 1. A piecewise-linear knotoid diagram is a union of finitely many edges: $[p_1, p_2], ..., [p_{n-1}, p_n]$ such that each edge intersects one or two other edges at the vertices, p_i , for i = 2, ..., n - 1. The vertices p_1 and p_n correspond to the endpoints of the diagram. Two edges can also intersect transversely at double points endowed with over/under-data, called crossings of the diagram.

Any classical knot diagram can be turned into a piecewise-linear knot diagram and piecewise-linear isotopy classes are in bijection with ambient isotopy classes of knots [12]. In a similar way any knotoid diagram can be turned into a piecewiselinear knotoid diagram. In this paper, we will be working for convenience with piecewise-linear knotoid diagrams. Moreover we will be considering $\Sigma = \mathbb{R}^2$.

2.1. Moves on knotoid diagrams. A Δ -move on a knotoid diagram is a replacement of an arc with two arcs (or vice versa) forming a triangle which does not contain any of the endpoints of the knotoid diagram and passing entirely over or under any arcs intersecting this triangle, as shown in Figure 2. The Reidemeister moves Ω_1 , Ω_2 , Ω_3 (Figure 2) are some special cases of Δ -moves. We shall call a Δ -move that takes place in a triangular region that does not contain any arcs of the diagram in its interior a planar Δ -move and denote it as Ω_0 -move. Finally, we have the swing moves whereby an arc containing an endpoint sweeps a triangular region free of any other arcs of the diagram. The swing moves can be viewed as special cases of Ω_0 -moves where one side of the isotopy triangle is omitted. See Figure 2. The moves consisting of pulling the strand adjacent to an endpoint over or under a transversal strand as shown in Figure 3 are the forbidden knotoid moves. It is clear that if both forbidden moves were allowed, any knotoid diagram in S^2 and in \mathbb{R}^2 could be turned into the trivial knotoid diagram.

Note. There are two situations where forbidden moves seemingly occur. Precisely, when the arc adjacent to an endpoint is involved in an Ω_1 - or an Ω_2 -move followed by a planar isotopy, as illustrated in Figure 4. These moves have the same effect as one or two consecutive forbidden moves of same type. We shall call these moves *fake forbidden moves*.



FIGURE 2. Δ -moves



FIGURE 3. Forbidden knotoid moves



FIGURE 4. Fake forbidden moves

We shall call $\Omega_0, \Omega_1, \Omega_2, \Omega_3$ moves together with the swing moves the Ω -moves. Two knotoid diagrams are *isotopic* to each other if there is a finite sequence of Ω -moves that transforms one into the other. The isotopy generated is clearly an equivalence relation and the isotopy classes of knotoid diagrams are called *knotoids* in \mathbb{R}^2 . The set of all knotoids in \mathbb{R}^2 is denoted by $\mathcal{K}(\mathbb{R}^2)$.

2.2. Extending the definition of knotoids. A multi-knotoid diagram in S^2 or in \mathbb{R}^2 [37] is an extended knotoid diagram having a finite number of knot diagrams in the sense that it is a generic immersion of the oriented unit interval and a finite number of oriented circles in S^2 or in \mathbb{R}^2 . The equivalence relation defined on knotoid diagrams applies to multi-knotoid diagrams directly, and the corresponding equivalence classes are called *multi-knotoids*. In this paper we work with multi-knotoids in \mathbb{R}^2 .

3. Braidoids

In this section we define braidoid diagrams and the isotopy classes of them that we call braidoids. A braidoid diagram is defined similarly to a braid diagram as a system of finite descending strands. The main difference is that a braidoid diagram has one or two of its strands starting with/terminating at an endpoint that is not necessarily at top or bottom lines of the defining region of the diagram.

3.1. The definition of a braidoid diagram.

Definition 2. Let I denote the unit interval $[0,1] \subset \mathbb{R}$. A braidoid diagram B is a system of a finite number of arcs immersed in $I \times I \subset \mathbb{R}^2$. We identify \mathbb{R}^2 with the *xt*-plane with the *t*-axis directed downward. The arcs of B are called the *strands* of B. Following the natural orientation of I, each strand is naturally oriented downward, with no local maxima or minima. There are only finitely many intersection points among the strands, which are transversal double points endowed with over/under data. Such intersection points are called *crossings* of B.

A braidoid diagram has two types of strands, the classical strands and the free strands. A *classical strand* is like a braid strand connecting a point on $I \times \{0\}$ to a point on $I \times \{1\}$. A *free strand* either connects a point in $I \times \{0\}$ or $I \times \{1\}$ to a special point that is located anywhere in $I \times I$ or connects two special points located

anywhere in $I \times I$. These special points are called the *endpoints* of *B*. A braidoid diagram contains either one free strand (see for example Figure 5(a)) or two free strands (see for example Figure 5 (b), (c), (d), (e)) and exactly two endpoints. The endpoints are specifically named as the *leg* and the *head* and are emphasized by graphical nodes labeled by *l* and *h*, respectively, in analogy with the endpoints of a knotoid diagram. Precisely, the head is the endpoint that is terminal for a free strand with respect to the orientation, while the leg is the starting endpoint for a free strand with respect to the orientation. See some examples of braidoid diagrams in Figure 5.



FIGURE 5. Some examples of braidoid diagrams

The ends of the strands of B other than the endpoints are called *braidoid ends*. We assume that braidoid ends lie equidistantly on the top and the bottom lines and none of them is vertically aligned with any of the endpoints. It is clear that the number of braidoid ends that lie on the top line is equal to the number of braidoid ends that lie on the bottom line of the diagram. The braidoid ends on top and bottom lines are arranged in pairs so that they are vertically aligned and are called *corresponding ends*. We number them with nonzero integers according to their horizontal order (from left to right), as in the examples illustrated in Figure 5.

Note that the endpoints of B differ conceptually from its braidoid ends. As we will see in the next section, the endpoints are subject to some isotopy moves unlike the braidoid ends, and the endpoints do not participate in the closure operation we introduce unlike the braidoid ends.

A braidoid diagram is *piecewise-linear* if all of its strands are formed by consecutive linear segments. Any braid diagram can be represented by a piecewise-linear braid diagram. Likewise, any braidoid diagram can be represented by a piecewiselinear braidoid diagram. We shall consider piecewise-linear braidoid diagrams, when convenient.

3.2. Braidoid isotopy. There are two types of local moves generating the braidoid isotopy.

3.2.1. Moves on segments of strands. We adapt the Δ -moves introduced in Section 2.1 to braidoid diagrams. A braidoid Δ -move replaces a segment of a strand with two segments in a triangular disk free of endpoints, passing only over or under the arcs intersecting the triangular region of the move whilst the downward orientation of the strands is preserved (see Figure 6). The oriented Ω_0 , Ω_2 and Ω_3 , which keep the arcs in the move patterns directed downward, can be viewed as special cases of braidoid Δ -moves.





FIGURE 6. A planar Δ -move on a braidoid diagram

3.2.2. Moves of endpoints. Like for knotoid diagrams, we forbid to pull/push an endpoint of a braidoid diagram over or under a strand, as shown in Figure 7. These are *forbidden moves* on braidoid diagrams. It is clear that allowing both forbidden moves can cancel any braiding of the free strands.



FIGURE 7. Forbidden braidoid moves

We allow the following moves on segments of braidoid strands containing endpoints.

- (1) Vertical Moves: As shown in Figure 8, the endpoints of a braidoid diagram can be pulled up or down in the vertical direction as long as they do not violate any of the forbidden moves (e.g. crossing through or intersecting any strand of the diagram). Such moves are called *vertical moves*.
- (2) Swing Moves: An endpoint can also swing to the right or the left like a pendulum (see Figure 9) as long as the downward orientation on the moving arc is preserved, and the forbidden moves are not violated.



FIGURE 8. A vertical move on h



FIGURE 9. The swing moves for braidoids

Definition 3 (braidoid isotopy). It is clear that assuming braidoid ends fixed at the top and bottom lines, the braidoid Ω_2 and Ω_3 - moves together with braidoid Ω_0 -moves and the swing and vertical moves for the endpoints generate an equivalence relation on braidoid diagrams in \mathbb{R}^2 . Two braidoid diagrams are said to be *isotopic* if one can be obtained from the other by a finite sequence of braidoid isotopy moves. An isotopy class of braidoid diagrams is called a *braidoid*.

4. FROM BRAIDOIDS TO PLANAR KNOTOIDS - A CLOSURE OPERATION

We present a closure operation on braidoid diagrams in analogy with the closure of braids in handlebodies [33]. In order to do this, we introduce the notion of labeled braidoids.

Definition 4. A labeled braidoid diagram is a braidoid diagram whose corresponding ends are labeled either with o or u in pairs. See Figures 10, 13. Each label indicates either an *overpassing* or *underpassing* arc, respectively, that will take place in the closure operation explained below.

Definition 5. Let *B* be a labeled braidoid diagram. The *closure* of *B*, denoted \hat{B} , is a planar (multi)-knotoid diagram obtained by the following topological operation: each pair of corresponding ends of *B* is connected with an embedded arc (with slightly tilted extremes) that runs along the right hand-side of the vertical line passing through the ends and in a distance arbitrarily close to this line. The connecting arc goes *entirely over* or *entirely under* the rest of the diagram according to the label of the ends. We demonstrate abstract and concrete examples in Figure 10 and 13, respectively.



FIGURE 10. The closure of an abstract labeled braidoid diagram

The reason that a joining arc is required to lie in an arbitrarily close distance to the line of the related corresponding ends is that, otherwise, forbidden moves may obstruct an isotopy of \hat{B} between any two joining arcs. Notice also that the resulting multi-knotoid depends on the labeling of the braidoid ends. In Figure 13 we see two labeled braidoid diagrams induced by the same underlying braidoid diagram via different labelings, give rise to non-equivalent knotoids.

On labeled braidoid diagrams we allow the braidoid Δ -moves and the vertical moves to take place on labeled braidoid diagrams and we forbid the forbidden braidoid moves. However, we do not allow swing moves for labeled braidoid diagrams in full generality. We only allow the *restricted swing moves* whereby the swinging of an endpoint takes place within the interior of the vertical strip determined by the neighboring vertical lines passing through two consecutive pairs of

corresponding braidoid ends (see Figure 11). The reason for restricting the swing moves is because if the endpoints surpass the vertical lines of the corresponding ends this will cause forbidden moves on the closure. See Figure 12 for an example.

Definition 6. Labeled braidoid isotopy is generated by the braidoid Ω -moves, the vertical moves and the restricted swing moves, preserving at the same time the labeling. Equivalence classes of labeled braidoid diagrams under this isotopy relation are called *labeled braidoids*.



FIGURE 11. The restricted swing moves for braidoids



FIGURE 12. A swing move causing a forbidden move on the closure

Lemma 1. The closure operation induces a well-defined mapping from the set of labeled braidoids to the set of multi-knotoids in \mathbb{R}^2 .

Proof. Let b_1 and b_2 be two labeled braidoid diagrams representing the same labeled braidoid. Then b_1 and b_2 differ from each other by braidoid isotopy moves. It is clear that braided Ω_2 and Ω_3 -moves are transformed into a sequence of knotoid Ω_2 and Ω_3 -moves. Also, the vertical and swing moves are transformed into planar isotopy on the (multi)-knotoid diagram obtained by the closure. Therefore the closures of b_1 and b_2 are isotopic (multi)-knotoid diagrams.



FIGURE 13. An example of non-equivalent labeled closures

5. An algorithm for obtaining labeled braidoids from planar knotoids

In this section we present the braidoiding moves on knotoid diagrams that induce algorithms for turning any planar (multi)-knotoid diagram into a labeled braidoid diagram [19]. By these algorithms we obtain the following theorem.

Theorem 1. (An analogue of the Alexander theorem for knotoids) Any (multi)knotoid diagram in \mathbb{R}^2 is isotopic to the closure of a labeled braidoid diagram.

In [19] two such algorithms for proving Theorem 1 are presented. One of the algorithms appearing also in [18] is conceptually lighter than the other one presented here, see also Remark 1. The algorithm we present here is more 'rigid' in the sense that it assumes knotoid diagrams as rigid diagrams. This makes the algorithm more appropriate for proving a braidoid equivalence result analogous to the classical Markov theorem for classical braids.

5.1. The basics of the braidoiding algorithm. Let K be a (multi)-knotoid diagram in a plane identified with the xt-plane. We describe below how to manipulate K in order to obtain a labeled braidoid diagram, after endowing the plane of K with top-to-bottom direction. We will be assuming that K lies in $[0, 1] \times [0, 1]$ since K is compact.

5.1.1. Up-arcs and free up-arcs. It is clear that by small perturbations K can be assumed to be a diagram without any horizontal or vertical arcs. Thus K consists of a finite number of arcs oriented either upward or downward, and these arcs are

separated by finitely many local maxima or minima. The arcs of K that are oriented upward are called *up-arcs* and the ones oriented downward are called *down-arcs* of K. An up-arc may contain crossings of different types (over/under-crossings) or no crossing at all. See Figure 14. An up-arc that contains no crossing is called a *free up-arc*.



FIGURE 14. Two up-arcs containing crossings and a free up-arc

5.1.2. Subdivision. We start by marking the local maxima and minima of K with points, which we name as subdividing points. In the process we may need to subdivide further some of the up-arcs of K so that each one contains crossings of only one type. We attach a label to each up-arc accordingly to the crossing type it contains: we attach o if the up-arc contains over-crossing(s), u if the up-arc contains under-crossing(s). The up-arcs that are free of crossings can be labeled either o or u.

5.1.3. Braidoiding moves. The basic idea of turning K into a labeled braidoid diagram is to keep the arcs of K that are oriented downward, with respect to the top-to-bottom direction, and to eliminate its up-arcs by turning them into braidoid strands. The elimination of the up-arcs is done by utilizing the sequence of moves illustrated in Figure 15. Precisely, a braidoiding move consists of cutting an up-arc at a point and pulling the resulting sub-arcs to top and bottom lines entirely over or entirely under the rest of the diagram by preserving the alignment with the cut-point. Finally we slide the resulting sub-arcs down and up, respectively, across local triangular regions in order to eliminate the upward oriented pieces. An up-arc is eventually turned into a braidoid strand as also depicted in Figure 15. It can also be verified by Figure 15 that the resulting ends obtained by cutting the up-arc QP at a point are pulled entirely over the rest of the diagram and received the label o, and when we join the resulting pair of corresponding ends with an over-passing arc, we obtain a closed strand that is isotopic to the initial up-arc QP.



FIGURE 15. A braidoiding move

5.1.4. Cut-points and sliding triangles. Let QP be an up-arc of K with respect to a given subdivision of K where Q denotes the initial and P denotes the top-most subdivision point.

The right angled triangle which lies below QP and admits QP as hypotenuse, and is a special case of a triangle enabling a Δ -move is called the *sliding triangle* of the up-arc QP. We denote the sliding triangle by T(P), see Figure 16. The disk bounded by the sliding triangle T(P) is utilized after cutting QP, for sliding down the resulting lower sub-arc across it. See Figure 16.



FIGURE 16. The sliding triangle of the up-arc QP with P the cut point

A *cut-point* of an up-arc is defined to be the point where the up-arc is cut to start a braidoiding move. We pick the top-most point $P \in QP$ as the cut-point of QP for our algorithm.

5.1.5. A condition on sliding triangles. One may come across the situation where one (or two) of the endpoints lies in the region bounded by some sliding triangle as in Figure 17. This is an unwanted situation since it would cause a forbidden move when the knotoid strand obtained by closing the resulting braidoid strand is tried to be isotoped back to the initial up-arc. We impose the following condition on knotoid diagrams to avoid this.

The endpoint triangle condition A sliding triangle of a knotoid diagram is not allowed to contain an endpoint.

To satisfy this condition we introduce a subdivision of up-arcs into smaller subarcs by adding extra subdividing points. See Figure 17. More precisely, we have the following proposition.

Lemma 2. Let K be a knotoid diagram and T(P) be the sliding triangle corresponding to an up-arc QP. If T(P) contains the leg or the head of K in its interior or boundary then there is a further subdivision of QP admitting new sliding triangles whose disks are disjoint from the endpoint.

Proof. We can assume that QP lies in the first quadrant of the plane and has a positive slope without loss of generality. There is a unique horizontal and vertical line passing through the endpoint in question and each intersecting QP exactly at one point, since $x_Q \leq x_{endpoint} \leq x_P$ and $t_P \leq t_{endpoint} \leq t_Q$. We pick an interior point in the small line segment whose boundary is the union of the two intersection points and declare it as a new subdividing point on QP. Let P^* denote the chosen

point. The sliding triangles intersecting at the point P^* are clearly smaller than the one with the top vertex P and they do not contain the endpoint. In case that T(P) contains two of the endpoints, we introduce two new subdividing points on QP each chosen as above and none of the corresponding sliding triangles contains the endpoints.



FIGURE 17. A further subdivision of the up-arc QP

5.2. Braidoiding algorithm. We now present our algorithm for obtaining a labeled braidoid diagram from a (multi)-knotoid diagram K. The algorithm runs as follows.

Step 1: Preparation for eliminating the up-arcs

- (1) The diagram is marked with subdividing points as explained in Section 5.1.2 so that the endpoint triangle condition is satisfied. It is furthermore assumed to satisfy some general positioning conditions, which can be ensured by 'small' isotopy moves, namely
- (2) no arcs is vertical or horizontal,
- (3) no subdividing points are vertically aligned with each other unless they share a common edge and neither with the endpoints or with any of the crossings,
- (4) the endpoints of K are assumed to lie on arcs that are directed downward.

Step 2: Applying the braidoiding moves

We order the up-arc and finally apply the braidoiding moves to each up-arc of K in the given order.

Figure 18 illustrates for the braidoiding algorithm with a concrete example.



FIGURE 18. An illustration for the algorithm

5.2.1. Obstructions for the braidoiding algorithm and resolutions. Now, we discuss some bugs of the braidoiding algorithm. In some cases, as exemplified in Figure 19, the algorithm is obstructed by a clasp occuring in the sliding triangle of an uparc. We see in the figure that the braidoiding move applied on the second-ordered up-arcs labeled with o and u respectively, cannot be completed, due to the clasps in their sliding triangles. It can be verified by checking all possible positionings, labelings and orderings of any two up-arcs that this type of obstruction may occur only if:

- the top-most point of the first-ordered up-arc intersects the sliding triangle of the second-ordered up-arc and,
- two up-arcs with intersecting sliding triangles, are labeled the same.





FIGURE 19. Obstructions for applying braidoiding moves

5.2.2. *Resolutions of the obstructions*. Due to the conditions creating obstructions, the resolutions for them can be:

- swapping the ordering of the up-arcs; see Figure 20,
- changing the label of the free up-arc if there is a free up-arc involved in an obstruction; see Figure 21,
- subdividing the up-arcs further to have disjoint sliding triangles; see Figure 22.



FIGURE 20. Swapping the order of up-arcs repairs the obstruction



FIGURE 21. Changing the label of the free up-arc repairs the obstruction



FIGURE 22. Adding a subdividing point to the upper up-arc repairs the obstruction

5.2.3. The classical triangle condition. We can make the braidoiding algorithm a simultaneous algorithm that is processed independently of the ordering of the uparcs, by imposing the following condition.

Definition 7. Two sliding triangles are said to be *adjacent* if the corresponding up-arcs have a common subdidiving point, and *non-adjacent* otherwise.

The *classical triangle condition* says that non-adjacent sliding triangles are allowed to intersect only if the up-arcs of the triangles have different labels. The classical triangle condition can always be satisfied by the following lemma.



FIGURE 23. The classical triangle condition

Lemma 3. Let K be a knotoid diagram. There exists a subdivision of K satisfying both the classical triangle condition and the endpoint triangle condition.

Proof. This lemma is proved similarly with Lemma 1 in [31]. For completeness, we adapt the proof here for the knotoid case. Let d_1 be the minimum distance between any two crossings appearing on the up-arcs of K. Choose some $r, 0 < r < d_1$ so that the disk of radius r centered at a crossing contained in an up-arc, intersects the up-arc at only four local strands around the crossing. Let d_2 be the minimum distance between any two disjoint points that are located outside the disks of radius r around the crossings and on different segments. Letting $0 < \epsilon < \frac{1}{2}min\{d_1, d_2\}$ be the distance between any two subdivision points on K provides a subdivision of K satisfying the classical triangle condition and also the triangle condition for the endpoints. Each sub-arc of length ϵ is also labeled according to the crossing type it contains, or if some become free, they are labeled freely.

5.2.4. Proof of Theorem 1. With the discussion above we know that there is always a choice for ordering and labeling of up-arcs of a (multi)-knotoid diagram and, moreover that we can impose the classical triangle condition to make the algorithm free of ordering. This is equivalent to saying that the braidoiding moves can be done all simultaneously and it is sufficient for ensuring that the algorithm terminates in a finite number of steps and always results in a labeled braidoid diagram. Finally, it is clear from Figure 15 that if we apply closure to the corresponding strands according to their label, then each pair can be contracted back to the initial up-arc. Such contractions utilize the isotopy moves for knotoids, thus the closure of the resulting braidoid diagram is isotopic to the (multi)-knotoid diagram we started with. QED

Remark 1. In [18] we present another braidoiding algorithm which is also induced by the same braidoiding moves. The main difference of that algorithm from the one we describe in this paper is that it firstly eliminates each up-arc of a knotoid diagram that contains a crossing by a 90-degree or 180-degree rotation of the crossing. The braidoiding moves are applied to the resulting up-arcs that are all free up-arcs. Furthermore, since the labels of the free up-arcs are not forced they can be labeled only u. Applying the braidoiding algorithm on such a knotoid diagram generates a labeled braidoid diagram whose strands are all labeled u. This induces the idea of closing braidoid diagrams in a uniform way, that is, by using only underpassing arcs that run in arbitrarily close distance to the vertical lines of pairs of corresponding ends, for connecting them. We call this closure the *uniform closure*.

The above remark leads to the following result.

Theorem 2. [19, 18] Any multi-knotoid diagram in \mathbb{R}^2 is isotopic to the uniform closure of a braidoid diagram.

6. L-Equivalence of braidoid diagrams

The L-moves were originally defined for classical braid diagrams [30, 31, 32] by the second listed author, and they were used for proving a one-move analogue of the classical two-move Markov theorem [34, 39, 9, 8, 35, 36, 10]. In the sequel, we adapt the L-moves for braidoid diagrams and use them for formulating a geometric analogue of the classical Markov theorem.

6.1. The L-moves.

Definition 8. An *L*-move on a labeled braidoid diagram B is the following operation:

- (1) Cut a strand of B at an interior point which is not vertically aligned with a braidoid end, an endpoint or a crossing of B. The existence of such point can be ensured by a general positioning argument.
- (2) Pull the resulting ends away from the cut-point to top and bottom respectively, keeping them vertically aligned with the cut-point, so as to create a new pair of braidoid strands with corresponding ends. The new strands run both entirely it over or *under* the rest of the braidoid diagram depending on the type of the *L*-move applied. There are two types of *L*-moves, namely L_{over} and L_{under} -moves, denoted by L_o and L_u respectively. An L_o -move comprises pulling the resulting sub-strands entirely over the rest of the diagram. An L_u -move comprises pulling the sub-strands entirely under the rest of the diagram. See top row of Figure 24.
- (3) After an *L*-move applied on a labeled braidoid diagram, the new pair of corresponding strands gets the labeling of the *L*-move: If the strands are obtained by an L_o -move then they are labeled o and if they are obtained by an L_u -move then they are labeled u. Then, as can be verified by Figure 24, the closure of a pair of labeled braidoid strands resulting from an *L*-move is isotopic to the initial arc.



FIGURE 24. L-moves and the closures of the resulting strands

18

In order to formulate a braidoid equivalence in analogy to the classical braid equivalence utilized in Markov theorem we also need to discuss the fake forbidden moves.

Definition 9. We define a *fake forbidden move* on a labeled braidoid diagram B to be a forbidden move on B which upon closure induces a sequence of fake forbidden moves on the resulting (multi-)knotoid diagram. A *fake swing move* is a swing move which is not restricted, in the sense that the endpoint surpasses the vertical line of a pair of corresponding ends but in the closure it gives rise to a sequence of swing and fake forbidden moves on the resulting (multi-)knotoid diagram. See Figure 25 for an example of a fake swing move and a fake forbidden move on a labeled braidoid diagram.



FIGURE 25. A fake swing and a fake forbidden move

Definition 10. The *L*-moves together with labeled braidoid isotopy moves and fake swing moves generate an equivalence relation on labeled braidoid diagrams that is called the *L*-equivalence. The *L*-equivalence is denoted by \sim_L .

The labeled braidoid that is associated to a knotoid K via a braidoiding algorithm, is not unique up to the braidoid isotopy. Precisely the labeled braidoid depends on the choices made for bringing the knotoid diagram to satisfy the general position requirements before starting the braidoiding algorithm, such as: arrangement of the endpoints, subdivision chosen on the arcs of the knotoid diagram and the labeling of the free up-arcs. Moreover, the knotoid diagram is subject to knotoid isotopy moves. We shall show that the labeled braidoid diagrams that are associated to K, are independent of algorithmic choices and isotopy moves up to L-equivalence.

Before stating our theorem we give the following definition.

Definition 11. A (multi)-knotoid diagram K in the xt-plane, with a subdivision and labeling on its up-arcs, is said to be in *general position* if

- it has no vertical or horizontal arcs,
- no subdividing point or endpoint is vertically aligned with a subdividing point, a crossing or an endpoint,
- the arcs adjacent to the endpoints of K are down-arcs,
- no sliding triangle encloses an endpoint,
- sliding triangles satisfy the classical triangle condition.

Note that a (multi)-knotoid diagram can be always brought to general position by small Δ -moves.

Theorem 3. [19] (An analogue of the Markov theorem for braidoids) The closures of two labeled braidoid diagrams are isotopic (multi)-knotoids in \mathbb{R}^2 if and only if the labeled braidoid diagrams are related to each other via L-equivalence moves.

Proof. For the proof of Theorem 3, we assume that (multi)-knotoid diagrams are in general position.

The 'if' part is clear from the definitions of L-moves and fake swing moves. More precisely, let B_1, B_2 be two labeled braidoid diagrams related to each other by an Lmove. Let the arc, illustrated in Figure 24, be a segment of a strand of B_1 on which an L-move is applied. It can be observed by the same figure that the closure of the resulting strands labeled accordingly to the type of the L-move applied, is isotopic to this arc. This implies that the closures of B_1 and B_2 are isotopic. From this, the closure map extends to a well-defined map cl_L on the set of all L-equivalence classes of labeled braidoids.

 $cl_L: \{L\text{-classes of labeled braidoid diagrams}\} \rightarrow \{\text{Multi-knotoids in } \mathbb{R}^2\}.$

For showing the 'only if' part, we need to show that the map cl_L is a bijection. We adapt these parts here and check the cases involving the endpoints. For this we first show the braidoiding algorithm induces a well-defined mapping br,

br: {Multi-knotoids in \mathbb{R}^2 } \rightarrow {L-classes of labeled braidoid diagrams},

that associates a (multi)-knotoid K in \mathbb{R}^2 to the *L*-class of the braidoid diagram obtained from any representative of K by the braidoiding algorithm. We call this map the *braidoiding map*.

In order to show the mapping br is well-defined we need to show that, up to *L*-equivalence, the resulting labeled braidoid diagram is independent of: the choices made for bringing a knotoid diagram into general position, choices of subdividing points for the up-arcs, labelings of free up-arcs, the Ω -moves of knotoid diagrams. The parts of the proof of Theorem 3 not involving the endpoints are analogous to the case of classical knots and braids [30, 31].

Lemma 4. Let K be a (multi)-knotoid diagram in \mathbb{R}^2 with a subdivision that satisfies the triangle conditions. Adding a subdividing point to an up-arc of K and labeling the new up-arcs same as the initial labeling yields an L-equivalent labeled braidoid diagram.

Proof. We first note that addition of a subdividing point does not violate the triangle conditions. Let QP denote an up-arc of K as depicted in Figure 26. First let us assume that the vertical line passing through an endpoint of K does not intersect QP. In this case, we can add a new subdividing point on QP and apply braidoiding moves at that point and the point P. The proof for showing that the resulting braidoid diagram is L-equivalent to the one obtained by applying a braidoiding move at the top-most point P follows similarly as for classical braid diagrams. The reader is directed to [31] for details.

Now let us assume that the vertical line passing through an endpoint of K intersects with QP at some point on QP. Then QP can be seen as the union of two sub-arcs joined at the intersection point and one of which contains the point P. The sub-arcs can be re-labeled accordingly to the initial labeling. By the classical

argument discussed above, if a new subdividing point is added on QP on the subarc containing P, then applying a braidoiding move at this point results in a labeled braidoid diagram that is L-equivalent to the labeled braidoid diagram obtained by applying a braidoiding move at the point P. Suppose now that a new subdividing point P_1 is chosen on the sub-arc that does not contain the point P. As we can verify in Figure 26, the labeled braidoid diagram resulting from a braidoiding move applied at the point P can be taken to the braidoid diagram resulting from the latter subdivision (that is by applying braidoiding moves at the points P and P_1) by an L-move that is applied at the point Q^* . Here the point Q^* denotes the intersection point of the vertical line passing through P_1 and the lower strand containing Q that has been obtained by the initial braidoiding of QP. Note that a neighborhood of Q containing the point Q^* is perturbed to slope slightly downwards for enabling application of the L-move.



FIGURE 26. Adding a subdividing point on the up-arc yields L-equivalence

Lemma 5. Labeling a free up-arc either with o or u does not change the resulting labeled braidoid diagram up to L-equivalence.

Proof. The proof is illustrated in Figure 27 and it goes in analogy with the classical case. In the figure, a free up-arc QP is labeled both with u and o. For simplicity we first assume that the sliding triangle of QP is also free of any arcs.

We start by applying a braidoiding move on QP labeled with u at the point P to obtain a pair of braidoid strands. Then we apply an L_o -move at a point (the point P^* in the figure) that is arbitrarily close to P so that there is no endpoint or

braidoid end in the vertical strip defined by P and P^* . Let Q^* be the point where the vertical line passing at P intersects the resulting lower piece of the sub-strand containing Q. We assume that the piece of strand containing Q^* slopes downward. Deletion of an L_u -move at the point Q^* cancels the pair of braidoid strands labeled u. By the same figure we can verify that the resulting labeled braidoid diagram can be turned into to the labeled braidoid diagram obtained by a braidoiding move on QP when it is labeled with o, by a sequence of L-moves: First by applying an L_o -move at Q^* and then deleting an L_o -move to cancel one of the pairs of braidoid strands.

The sliding triangle of the up-arc may intersect with other arcs of the diagram as illustrated in Figure 28. In this figure, we show a case where a free up-arc labeled with u intersects the sliding triangle of the free up-arc labeled with o. By using Lemma 4, we first subdivide the free up-arc labeled with o into small enough subarcs to make all disks of the corresponding sliding triangles free of arcs. We now re-label each of the free up-arcs resulting from the new subdivision with o. By the discussion above, we know that the diagram obtained by applying braidoiding moves at each added subdividing points on the up-arc is L-equivalent to the one obtained by applying braidoiding moves to the up-arc in the third row where each sub-arc is labeled with u. Finally, again by Lemma 4 we know that if we delete all the added subdividing points on the up-arc keeping the initial subdividing points and the label u then by a braidoiding move we obtain a labeled braidoid diagram that is L-equivalent to the initial labeled braidoid diagram obtained by a braidoiding move applied on the up-arc labeled with u. Notice that, to ensure the classical triangle condition in the final step we also change the labeling of the intersecting arc from u to o.



FIGURE 27. Different labels on a free up-arc





FIGURE 28. Re-labeling the free up-arcs

Corollary 1. If we have an appropriate choice of relabelling the up-arcs resulting from a further subdivision on K then by Lemmas 4 and 5, the resulting labeled braidoid diagrams are L-equivalent.

Lemma 6. Two labeled braidoid diagrams that are obtained with respect to any two subdivisions S_1, S_2 on a knotoid diagram K that satisfy the triangle conditions, are L-equivalent to each other.

Proof. It is clear that any subdivision further than S_1 or S_2 satisfies both triangle conditions. Consider the further subdivision $S_1 \cup S_2$ of both S_1 and S_2 on K. By Lemmas 4 and 5, the labeled braidoid diagram that results from the subdivision $S_1 \cup S_2$ is L-equivalent to the labeled braidoid diagrams that result from the subdivision S_1 and S_2 . Since the L-equivalence is an equivalence relation, the lemma follows.

Finally we shall show the *L*-equivalence under the Ω -moves.

Lemma 7. Two (multi-)knotoid diagrams in \mathbb{R}^2 are related to each other by planar isotopy and Ω - moves only if the corresponding labeled braidoid diagrams are related to each other by labeled braidoid isotopy moves, L-moves, fake swing moves or fake forbidden moves.

Proof. Let us start with the observation that by imposing the classical triangle condition, we can assume that the knotoid isotopy moves take place without interfering with the braidoiding process of the rest of the diagrams that lie outside the local regions of the moves. In fact we can assume the up-arcs outside the move disks are all turned into braidoid strands and only the arcs lying in the move disks are left for elimination.

Examining the Ω_1 , Ω_2 and Ω_3 -moves follows similarly with the examination of Reidemeister moves on classical knots/links under the braiding moves [32, 31]. Here in Figure 29 we give an illustration for the Ω_1 -move away from the endpoints and how it is transformed to an *L*-move under a braidoiding move.

23



FIGURE 29. An Ω_1 -move under braidoiding

In this setting, it is crucial to examine specifically the swing moves displacing the endpoints of a knotoid/multi-knotoid diagram. A swing move may displace an endpoint that lies on a down-arc and in a way that the endpoint does not change its position with respect to a vertical line passing through a cut-point in the diagram then the resulting (labeled) braidoid diagrams are clearly related to each other by a restricted swing move.

A swing move may move an endpoint so that the swinging arc changes from being a down-arc to being an up-arc by keeping the endpoint at the same vertical alignment as in Figure 30. In the braidoiding process, the down-arc is kept but a braidoiding move is applied on the resulting up-arc. It can be verified by the figure that the resulting labeled braidoid diagrams obtained from two knotoid diagrams which are related by such swing move, are related to each other by an *L*-move. See Figure 30. Note that the points P and P^* shown in the figure that are chosen for applying the L_o -move on the resulting braidoid diagram and the braidoiding move on the resulting knotoid diagram, respectively, are vertically aligned.

A swing move may also cause the endpoint to change its position with respect to a vertical line passing through a cut-point on the diagram as illustrated in Figure 31. This means that the endpoint of the swinging arc in the resulting labeled braidoid diagram crosses the vertical line passing through the new pair of corresponding braidoid ends. This situation can be examined in two separate cases. One case is that the swinging arc is a down-arc and remains as a down-arc during the move. In this case, no braidoiding move applies on the arc and the resulting braidoid diagrams are related to each other by a fake forbidden move (recall Definition 9). The other case is that the swinging arc is an up-arc and the endpoint is displaced with respect to a vertical line passing through a cut-point lying on some up-arc which can be the swinging arc itself. In this case, we apply a last braidoiding move on the swinging arc before and after the move. Then the comparison of the resulting (labeled) braidoid diagrams is reduced to the previous case and they are related to each other by a fake swing move when the cut-point lies on the swinging arc or a fake forbidden move as discussed in the previous case.

From the above discussion we can deduce Lemma 7.



FIGURE 30. A swing move changing a down-arc to an up-arc



FIGURE 31. A swing move displacing an endpoint with respect to a cut-point

Lemma 8. A fake forbidden move on a labeled braidoid diagram is generated by L-moves, planar isotopy moves and fake or restricted swing moves.

Proof. The proof can be verified by Figure 32. The figure illustrates a finite sequence of planar isotopy moves, addition/deletion of L_u -moves and restricted swing moves that gives the fake forbidden move. This can be varied to include L_o -moves (when the swinging arc goes under a braidoid strand) and fake swing moves (when

the swinging arc swings across the vertical line determined by the braidoid end that is connected to the swinging endpoint with the swinging arc, see the first instance of Figure 25. Figure 33 also illustrates the statement on a complete labeled braidoid diagram.



FIGURE 32. A fake forbidden move via L-moves and a swing move



FIGURE 33. A fake forbidden move on a labeled braidoid diagram

Lemma 9. Two (multi-)knotoid diagrams in \mathbb{R}^2 are related to each other by planar isotopy and Ω -moves only if the corresponding labeled braidoid diagrams are L-equivalent.

Proof. The lemma follows from Lemma 7 and Lemma 8.

26

From Lemma 6 and Lemma 9 it follows that the braidoing map br is a welldefined map. Lastly it needs to be shown that br is the inverse map of cl_L . Let B be a labeled braidoid diagram. It is clear that the closure diagram of B, \hat{B} , is a knotoid diagram in general position whose only up-arcs are the connection arcs taking place in closing B. The braidoiding algorithm eliminates each up-arc of \hat{B} and turns it into a labeled braidoid diagram which is isotopic to B. Thus,

$$br \circ cl_L = id$$

Given a knotoid diagram K in general position, by applying the braidoiding algorithm we obtain a labeled braidoid diagram B. Clearly the closure of any labeled braidoid diagram that is L-equivalent to B is isotopic to K. Hence it follows

$$cl_L \circ br = id.$$

By the above the proof of Theorem 3 is now completed.

7. From braidoids to braids

In this section we show another application of the *L*-moves, namely the induced relation between the set of all braidoid diagrams and the set of all classical or virtual braid diagrams.

We define a *virtual braidoid diagram* to be a braidoid containing classical as well as virtual crossings and a *virtual braidoid* to be the equivalence class of virtual braidoid diagrams under the moves of the braidoid diagrams extended by the virtual braid moves. For more on virtual braids and the virtual braid group the reader may consult [25, 26, 6] and references therein. Further, for analogues of the Alexander and the Markov theorems for virtual knots and virtual braids see [27, 23].

In analogy to knotoids closing to classical or virtual knots, a braidoid resp. a virtual braidoid diagram is closed to a classical resp. a virtual braid diagram by connecting its endpoints with a simple arc in the plane. There are only finitely many intersection points between the connecting arc and the braidoid diagram, which are transversal double points endowed with under, over resp. virtual crossing data. By a classical topological argument, the arc connecting the endpoints of a braidoid diagram is unique up to isotopy in the classical or in the virtual sense.

7.1. The underpass closure and the virtual closure. We fix the connecting arc to be passing *under* any other arc. In order to receive as outcome a braid diagram we must also ensure the braid monotonicity condition. There can be two cases for the connection. If the head of the braidoid diagram appears before its leg (as t increases) then a downward directed arc can be chosen to connect them, see the top row of Figure 34 for an abstract illustration. If, however, the leg of the braidoid diagram appears before its head then any arc connecting them is an arc that is directed upward, an up-arc, in the resulting tangle diagram. In this case, we apply a braiding move on the connecting arc which turns it into two new corresponding braid strands. See the bottom row of Figure 34 for an abstraction of this case.

Definition 12. For any given braidoid diagram we define the *underpass closure* of its endpoints as follows. If the head is in a relatively higher position than the leg, we join them together with a straight under-passing arc. If the leg is located higher than the head then we extend both endpoints by a pair of corresponding

underpassing braid strands, emanating from the leg and the head respectively, and vertically aligned with a point on the straight arc connecting the two endpoints.

The above apply analogously for the case where the connecting arc is passing *virtually* any other arc. In this case we obtain in the end a virtual braid. and call the closure the *virtual closure*. Especially in the case where the leg of the braidoid diagram appears before its head we eliminate the joining up-arc by a virtual braiding move, whereby all crossings in the resulting new pair of strands are virtual, see [27].



FIGURE 34. The underpass closure

7.2. The induced mappings. We shall now establish that the underpass closure (resp. the virtual closure) defined on braidoid diagrams induces a well-defined map on the set of braidoids (that is, isotopy classes of braidoid diagrams). We have the following proposition.

Proposition 1. The underpass closure is a well-defined surjective map from the set of braidoids to the set of L-equivalence classes of classical braids. Similarly, the virtual closure is a well-defined surjective map from the set of virtual braidoids to the set of virtual L-equivalence classes of virtual braids.

Proof. It is straightforward to show that the underpass closure is a surjective map. Indeed, by cutting out from a given braid diagram an open arc that either contains no crossings or it is an underpassing arc (resp. a virtual arc), we obtain a braidoid diagram whose underpass closure (resp. virtual closure) is clearly the original braidoid (resp. virtual braidoid) diagram.

Consider a braidoid isotopy on a given braidoid diagram (resp. virtual braidoid isotopy on a virtual braidoid diagram). The isotopy may keep the endpoints fixed

or it may change their relative vertical or horizontal positions. If the isotopy keeps the endpoints fixed then after the underpass (resp. virtual) closure this isotopy clearly transforms into braid isotopy (recall Section 3.2). Suppose that a swing move on one of the endpoints takes place. If the connecting arc is a down-arc then the corresponding braid diagrams clearly differ by braid isotopy (resp. virtual braid isotopy). If, however, the connecting arc is an up-arc then the swing move may change the vertical level (the ordering amongst the strands) of the resulting two braid strands. For an illustration see Figure 35. Yet, the two resulting braids will differ by conjugation, which is known to be a special case of *L*-equivalence [32] (resp. virtual *L*-equivalence [27]).

Suppose now that a vertical move on the endpoints takes place. If this move does not change the relative heights of the endpoints, it is clear that the connecting arcs are isotopic. If, however, the move changes the relative heights of the endpoints then the connecting arc changes from a down-arc to an up-arc or vice versa, see Figure 36. In this case, the two resulting braid diagrams differ by an L_{u} - move (resp. virtual *L*-move) applied on the connecting down-arc of the one diagram and at the same vertical level as that of the braiding move applied on the connecting up-arc of the other diagram.

By the arguments above the proof of Proposition 1 is completed.



FIGURE 35. A swing move transforms into an L-move on the underpass closure



FIGURE 36. A vertical move transforms into an L-move on the underpass closure

8. DISCUSSION

The theory of braidoids is a new diagrammatic setting extending the classical braid theory and giving rise to many new problems; the underlying algebraic structure for braidoids is not known yet and we will examine in a subsequent paper. However, we know how to partition a braidoid diagram into smaller diagrams that we call elementary blocks [18, 19]. Knots and knotoids have been used for topological tabulation of proteins which are open molecular chains. In [18] we suggest that braidoids can set up an algebraic way for tabulating proteins by the use of elementary blocks.

Acknowledgement

This research was partially supported through the programme 'Oberwolfach Leibniz Fellows' by the Mathematisches Forschungsinstitut Oberwolfach in 2019.

The first author cordially thanks the MFO for providing her a peaceful research environment during her stay as a Leibniz Fellow.

References

- Adams, C., Henrich, A., Kearney, K., and Scoville, N., Knots related by knotoids, Knots related by knotoids, *The American Mathematical Monthly*, 126:6, 483-490, DOI; 10.1080/00029890.2019.1583971
- [2] Alexander, J.W. A lemma on systems of knotted curves. Proc. Nat. Acad. Sci. U.S.A.(1923) 9, 93–95.
- [3] Artin E. Theorie der Zöpfe. Abh. Math. Sem. Hamburg Univ.(1926) 4, 47-72.
- [4] Artin, E. Theory of braids. Annals Math. (1947) 48, 101-126.
- [5] Barbensi, A., Buck, D., Harrington, H.A., and Lackenby M. Double branched covers of knotoids, arXiv:1811.09121 [math.GT] (2018).
- [6] Bardakov, V.G. The virtual and universal braids, Fundamenta Mathematicae 184 (2004) 159–186.
- [7] Bartholomew, A. Knotoids, A. Bartholomew's mathematical page, www.layer8.co.uk/maths.knotoids/index.htm, Jan. 2015.
- [8] Bennequin, D. Entrlacements et équations de Pfaffe, Asterisque 107-108 (1983) 87-161.
- [9] Birman J.S. Braids, links and mapping class groups. Annals of Mathematics Studies (1974) 82, Princeton University Press, Princeton.
- [10] Birman J.S., Menasco W.W. On Markov's Theorem. J. Knot Theory& Ramifications(2002)11 no.3, 295–310.
- [11] Brunn H. Über verknotete Curven. Verh. des intern. Math. Congr.(1897) 1, 256–259.
- [12] Burde, G., Zieschang, H. Knots, de Gruyter Stud. Math., vol. 5, Walter de Gruyter, New York, 2003, 559 pp.
- [13] Goundaroulis D., Dorier J., Benedetti F., Stasiak A. Studies of global and local entanglements of individual protein chains using the concept of knotoids. *Sci. Reports*(2017) 7, 6309.
- [14] Goundaroulis D., Gügümcü N., Lambropoulou S., Dorier J., Stasiak A., Kauffman L.H. Topological models for open knotted protein chains using the concepts of knotoids and bonded knotoids. *Polymers*, Special issue on Knotted and Catenated Polymers, Dusan Racko and Andrzej Stasiak Eds. 9(9) (2017), 444, DOI;10.3390/polym9090444.
- [15] Dabrowski-Tumanski, P., Rubach, P., Goundaroulis, D., Dorier, J., Sułkowski, P., Millett, K.C., Rawdon, E.J., Stasiak, A., and Sułkowska, J.I.KnotProt 2.0: a database of proteins with knots and other entangled structures, *Nucleic Acids Research* 1 (2018). DOI: 10.1093/nar/gky1140.
- [16] Daikoku, K., Sakai, K., and Takase, M.. On a move reducing the genus of a knot diagram, *Indiana University Mathematics Journal* 61.3 (2012), pp. 1111–1127. ISSN: 00222518, 19435258. url: http://www.jstor.org/stable/24904076.
- [17] Gügümcü, N., Kauffman, L.H. New invariants of knotoids. European J. of Combinatorics (2017), 65C, 186-229.
- [18] Gügümcü, N., Lambropoulou, S. Knotoids, Braidoids and Applications. Symmetry, Special Issue:Knot Theory and Its Applications (2017)
- [19] Gügümcü, N. On Knotoids, Braidoids and Their Applications. PhD thesis, 2017.
- [20] Gügümcü, N., Nelson, S., Biquandle coloring invariants of knotoids, J. Knot Theory & Ramifications, Vol. 28, No. 04, 1950029 (2019)
- [21] Jones, V.F.R., A polynomial invariant for knots via von Neumann algebras, Bull. Amer. Math. Soc. (N.S.) 12(1) (1985), 103–111.
- [22] Jones, V.F.R. Hecke algebra representations of braid groups and link polynomials, Ann. Math. 126 (1987), 335–388.
- [23] Kamada, S. Braid representation of virtual knots and welded knots, Osaka J. Math. 44 (2007), no. 2, 441–458. See also arXiv:math.GT/0008092.
- [24] Kassel, C., Turaev, V. Braid groups, Volume 247 of Graduate Texts in Mathematics, Springer, New York, 2008.
- [25] Kauffman, L.H. Virtual Knot Theory, European J. Comb. 20 (1999) 663-690.
- [26] Kauffman, L. H., Lambropoulou, S. Virtual braids. Fundamenta Mathematicae (2004), 184, 159-186.

- [27] Kauffman, L.H., Lambropoulou, S. Virtual braids and the L-move, J. Knot Theory & Ramif. 15(6) (2006), 773-811.
- [28] Kodokostas, D., Lambropoulou, S., Rail knotoids, to appear in J. Knot Theory & Ramif. (2019), arXiv:1812.09493.
- [29] Korablev, P.G., May, Y.K., Tarkaev, V., Classification of low complexity knotoids, Siberian Electronic Mathematical Reports (2018), 15, 1237-1244. [Russian, English abstract]. DOI: 10.17377/semi.2018.15.100.
- [30] Lambropoulou, S. Short proofs of Alexander's and Markov's theorems. Warwick preprint, 1990.
- [31] Lambropoulou, S. A study of braids in 3-manifolds. PhD thesis, University of Warwick, 1993.
- [32] Lambropoulou S., Rourke C. P. Markov's theorem in 3-manifolds. Topology and its Applications(1997), 78, 95-122.
- [33] Lambropoulou S., Häring-Oldenburg R., Knot theory in handlebodies, J. Knot Theory Ramif. (2002), 11, No. 6, 921-943, https://doi.org/10.1142/S0218216502002050.
- [34] Markov, A. A. Über die freie Äquivalenz geschlossener Zöpfe. Rec. Math. Moscou(1936), 1 (43), 73-78.
- [35] Morton, H. R. Threading knot diagrams. Mathematical Proceedings of the Cambridge Philosophical Society(1986), 99, 247-260.
- [36] Traczyk, P. A new proof of Markov's braid theorem. Banach Center Publications 1992, 42, Institute of Mathematics Polish Academy of Sciences, Warszawa (1998).
- [37] Turaev V. Knotoids. Osaka J. Math. 49 (2012), 195–223.
- [38] Vogel P. Representation of links by braid: A new algorithm. Commentarii Mathematici Helvetici (1990), 65, 104-113.
- [39] Weinberg N. Sur l' equivalence libre des tresses fermée. Comptes Rendus (Doklady) de l' Académie des Sciences de l' URSS (1939), 23(3), 215–216.
- [40] Yamada S. The minimal number of Seifert circles equals the braid index of a link. Invent. Math.(1987), 89, 347–356.

IZMIR INSTITUTE OF TECHNOLOGY DEPARTMENT OF MATHEMATICS GÜLBAHÇE CAMPUS URLA 35430 İZMIR, TURKEY AND UNIVERSITY OF GOETTINGEN MATHEMATICS INSTITUTE, BUNSEN-STRASSE 3-5 37073 GOETTINGEN, GERMANY

School of Applied Mathematical and Physical Sciences National Technical University of Athens Zografou Campus GR-15780 Athens, GREECE

Email address: sofia@math.ntua.gr

URL: http://www.math.ntua.gr/~sofia/

32